1 **REVISION 1**

2 **A new telluride topology: the crystal structure of honeaite Au₃TlTe₂.**

3 Mark D. Welch^{1*}, J.W. Still², C. M. Rice², C.J. Stanley¹

4 ¹Department of Earth Sciences, Natural History Museum, Cromwell Road, London SW7
5 5BD, United Kingdom.

6 ²Department of Geology & Petroleum Geology, School of Geosciences, Meston Building,
7 Kings College, Aberdeen AB24 3UE, United Kingdom

8 * mdw@nhm.ac.uk

9 Key words: honeaite, new mineral, crystal structure, telluride

10 **Abstract**

11 The crystal structure of the first thallium-bearing gold telluride, honeaite Au₃TlTe₂, is
12 reported and its topological novelty discussed. Honeaite is orthorhombic, space group *Pbcm*
13 and unit-cell parameters $a = 8.9671(4) \text{ \AA}$, $b = 8.8758(4) \text{ \AA}$, $c = 7.8419(5) \text{ \AA}$, $V = 624.14(6)$
14 $\text{\AA}^3$ ($Z = 4$). Its structure has been refined to $R_1 = 0.033$, $wR_2 = 0.053$, $GoF = 1.087$. The
15 structure is based upon a corrugated double-sheet comprising two sub-sheets, each composed
16 of six-membered rings of corner-linked TeAu₃ pyramids in which the Te lone-pair is
17 stereoactive. Rows of thallium atoms lie in the grooves between sheets and provide the only
18 inter-sheet connectivity *via* Tl-Au bonds. There is extensive Au-Au bonding linking the two
19 sub-sheets of the double-sheet. The structure is distinct from those of the 1:2 (Au,Ag)-
20 tellurides calaverite AuTe₂, sylvanite AuAgTe₄, and krennerite Au₃AgTe₈, which are based
21 upon sheet structures with no connecting inter-sheet atoms. It also differs fundamentally from
22 the structures of synthetic phases Ag₃TlTe₂ and Ag₁₈Tl₄Te₁₁, both of which have an

23 analogous stoichiometry. In contrast to the pyramidal TeAu_3 group of honeaite and
24 krennerite, Ag does not form the corresponding TeAg_3 group in its tellurides.

25

26 **Introduction**

27 Tellurium is one of the few elements to form stoichiometric compounds with gold in
28 nature. Recently, a new natural gold telluride was found at the Karonie gold deposit in the
29 Eastern Goldfields region of the Yilgarn Craton, Western Australia. The telluride has a novel
30 composition Au_3TlTe_2 and structure. There is no synthetic analogue. The mineral is named
31 honeaite after the mineralogist Russell M. Honea (1929-2002) who, amongst many diverse
32 achievements in geology, published the original description of the geology and mineralogy of
33 the Karonie deposit. The mineral and name have been approved by the International
34 Mineralogical Association as a new species (Rice et al. 2015). The geological occurrence,
35 petrogenesis and formal mineralogical description of honeaite are described in a separate
36 paper (Rice et al. in press).

37 It appears that a honeaite-like natural phase was found by Nechaev and Cook (2000)
38 at the Maiskoe gold deposit in Ukraine. Two microprobe analyses were reported that gave
39 compositions ($\text{Te} = 2 \text{ apfu}$) of $\text{Au}_{3.38}\text{Tl}_{1.14}\text{Te}_2\text{Sb}_{0.03}$ and $\text{Au}_{3.22}\text{Ag}_{0.02}\text{Tl}_{1.00}\text{Te}_2\text{Sb}_{0.03}$. No crystal
40 structure or unit-cell parameters were reported and no formal new-mineral description was
41 submitted to IMA or a name chosen. Consequently, while the composition is essentially that
42 of honeaite, it is unclear if it is actually this mineral. It would be of interest to obtain a
43 suitable sample of this material for structure determination to see if it is indeed honeaite.
44 Here, we report the determination of the structure of honeaite and discuss its topological
45 significance and distinctiveness.

46

47 **Sample material**

48 The crystal used in the structure determination reported below came from a rock core
49 sample collected at the Archaean Karonie gold deposit of the Eastern Goldfields province of
50 Western Australia. The geological occurrence of the new mineral honeaite is reported by Rice
51 et al. (in press). The type specimen and crystal are held in the collection of the Natural
52 History Museum (London) under catalogue number BM2015,36.

53 Chemical analyses of 17 grains of honeaite (one analysis per grain) in a polished thin
54 section were obtained, including for the type crystal, by electron microprobe analysis using
55 energy dispersive spectrometry operated at 15 kV, 2.8 nA, with a beam diameter of
56 approximately 0.005 mm and a live-time of 200 seconds. Element standards used were Au
57 metal (Au), thallium iodide (Tl), Te metal (Te). No other elements were detected in EDS
58 survey scans. Analytical totals were 99.60–101.02 %. The average composition of 17
59 analyses is 56.33(0.37) wt% Au, 19.68(0.27) wt% Tl and 24.30(0.11) wt% Te. The
60 corresponding average chemical formula is $\text{Au}_{2.996}\text{Tl}_{1.009}\text{Te}_{1.995}$. A spot analysis of the type
61 crystal gave 56.19 wt% Au, 19.45 wt% Tl and 24.19 wt% Te, with a corresponding chemical
62 formula of $\text{Au}_{3.003}\text{Tl}_{1.002}\text{Te}_{1.996}$.

63

64 **X-ray diffraction**

65 *Experimental methods and materials*

66 A polished petrographic thin section (0.03 mm thick) was surveyed by electron
67 microprobe analysis and numerous crystals of the new phase were identified as having a
68 distinctive new composition close to Au_3TlTe_2 . A small rectangular, flake-like crystal (0.040

69 $\times 0.035 \times 0.015$ mm) was extracted from the thin section by excavating with a fine
70 acupuncture needle (0.008 mm tip) and attached to a non-diffracting amorphous carbon fibre
71 (0.01 mm diameter) itself glued to a glass fibre support. Data were collected using an
72 XcaliburE single-crystal X-ray diffractometer equipped with an EoS CCD detector ([®]Rigaku
73 Oxford Diffraction) and graphite-monochromatised MoK α radiation operated at 45 kV, 40
74 mA. The data collection strategy used was chosen on the basis of a 2-hour pre-experiment. A
75 whole sphere of reflections was collected to 60° 2 θ . Information relating to the data collection
76 is given in **Table 1**. Raw reflection intensities were corrected for Lorentz and polarisation
77 effects and for absorption, the latter by the multi-scan method (averaging of equivalent
78 reflections) in the program *CrysAlisPro* ([®]Rigaku Oxford Diffraction) and converted to
79 structure factors.

80

81 **Results**

82 *Structure refinement*

83 The unit-cell of honeaite is metrically orthorhombic and systematic absences
84 indicated space group *Pbcm*. The structure was solved by Direct Methods and refined using
85 SHELX (Sheldrick, 2008). Neutral atomic scattering factors taken from *International Tables*
86 *for Crystallography* (Wilson, 1992) were used. Information relating to the data collection and
87 structure refinement is summarised in **Table 1**. Atom coordinates and displacement
88 parameters are given in **Table 2**; key interatomic distances are given in **Table 3**. A CIF file is
89 deposited with the journal.

90 The structure was solved and refined using SHELX (Sheldrick, 2008) within the
91 WinGX program suite (Farrugia, 1999). All six atoms of the asymmetric unit were found by

structure solution using Direct Methods. There are two Te, three Au and one Tl in the asymmetric unit; all atoms have site multiplicity 4. Assignment of Te versus (Au, Tl) used site scattering values (50 versus 79/81 electrons). Assignment of Au ($Z = 79$) versus Tl ($Z = 81$) was based upon crystal chemical reasoning: observed Te bond distances, all of which are almost identical (2.607–2.620 Å) and topological similarity with Te coordination in calaverite, AuTe_2 , strongly point to the presence of TeAu_3 groups. A Te–Au distance of 2.61 Å occurs in calaverite (Schütte and de Boer, 1988). The highly regular nature of the TeAu_3 group (bonds and angles) indicates that a single type of element coordinates Te in this group, i.e. Au. It cannot be TeTl_3 because this would require a much higher Tl content than the empirical formula has. Mixed Te(Tl,Au)_3 is unsupported by the bond lengths, all of which are almost the same and recognisable as Te–Au distances in related minerals calaverite, sylvanite and krennerite (Schütte and de Boer, 1988; Tunell, 1941; Dye and Smyth, 2012). Finally, Au–Au distances are 2.87 – 2.91 Å and are very similar to that of elemental gold (2.87 Å). Consequently, we assign all Au to the sites at 2.61 Å from Te and forming TeAu_3 groups. The maximum residual unmodelled electron density of $2 \text{ e}/\text{\AA}^3$ lies at 2.06 Å from Te(2) and is unassigned.

Structure topology

The structural features of honeaite are shown **Figures 1-4**. The structure comprises corrugated sheets of connected 6-membered rings of TeAu_3 groups $\parallel(100)$ as shown in **Figures 1 and 2**. Individual sheets comprise alternating chains of Te(1)Au_3 and Te(2)Au_3 pyramids. The overall structure is composed of stacked double-sheets in which Au–Au bonds connect the two sub-sheets (**Fig. 2**). Adjacent double-sheets are connected only *via* Tl–Au

bonds (2.966 Å). Rows of thallium atoms extend $\parallel[001]$ and lie in the grooves between sheet corrugations.

Thallium is coordinated to four Au atoms with bond lengths of 2.96–3.13 Å. There are also two very long distances to Te(1) atoms (3.49 Å), leading to an octahedral environment (**Fig. 3**). The exact nature of the Tl–Te distance is unclear, as the role of the Te lone-pair is undefined. The thallium site is very different from the three Au sites.

TeAu₃ groups

All Te and Au atoms form TeAu₃ groups in which Te is displaced to one side of the triangle of Au atoms and is characteristic of lone-pair stereoactivity. The two non-equivalent TeAu₃ groups are highly regular flattened triangular-based pyramids with Te–Au distances of 2.607–2.620 Å and Au–Te–Au angles of 97–103°. The polyhedral environments of Te(1)Au₃ and Te(2)Au₃ are very different, as illustrated in **Figure 4**. The Te(1)Au₃ pyramid shares two edges with TlAu₄ groups and apices with two Te(1)Au₃ and one Te(2)Au₃. The Te(2)Au₃ pyramid has no shared edges, but is corner-linked to two Te(2)Au₃, one Te(1)Au₃ and two TlAu₄.

Au polyhedra

The three Au sites (**Fig. 3**) have different coordination environments. Au(1) and Au(3) are octahedrally-coordinated, whereas Au(2) is four-fold coordinated. Au(1) is bonded to two Te, three Au atoms and one Tl atom; Au(3) is bonded to two Te, two Au and two Tl atoms; Au(2) is bonded to two Te atoms, one Au atom and one Tl atom. Au–Au distances are 2.87 – 2.91 Å and are very similar to that of elemental gold (2.87 Å). Au–Tl bonds are 2.97 – 3.06 Å. Te–Au bonds are all very similar at 2.61 – 2.62 Å. The connectivity of the Au

polyhedra is shown in **Figure 3**. The two octahedra (Au 1,3) share an edge and the four-coordinate Au(2) quadrilateral interpenetrates the Au(1) octahedron.

Thallium

As described above, the assignment of Tl *versus* Au sites can be made using crystal chemical reasoning. The four nearly coplanar Tl-Au bonds are long (2.97 – 3.13 Å) and uncharacteristic of Au-Au, one of which, Tl-Au(2), links adjacent double-sheets. Two very long distances (3.49 Å) to Te(1) atoms suggest octahedral coordination of Tl, although it is questionable if these two distances are meaningful bonds. TlAu₄ groups are corner-connected to form rows $\parallel[001]$.

Discussion

The synthetic compounds CsAu₃S₂, RbAu₃Se₂ and CsAu₃Se₂ (Klepp and Weithaler, 1996) have an analogous stoichiometry to that of honeaite, but the monovalent species is an alkali metal, whereas it is thallium in honeaite. The structure topology shared by these three compounds comprises alternating planar sheets of SeAu₃ (SAu₃) pyramids and Cs (Rb) atoms, and contrasts with the corrugated sheet of honeaite. Furthermore, the coordination of Se and S atoms is very different from Te in honeaite: Se and S are octahedrally-coordinated to three Au and three Cs (Rb) cations, e.g. Se[Au₃Cs₃] in CsAu₃Se₂.

One of the two non-equivalent Te atoms of honeaite has possible bonding distances to Tl, whereas the other Te is only bonded to Au. In both cases, Te[Au₃Tl₂] and Te[Au₃], the coordination is highly asymmetric (one-sided), suggesting stereoreactivity of both Te atoms. For CsAu₃Se₂ and CsAu₃S₂, the Cs-Se and Cs-S distances of 3.76 and 3.73 Å, respectively,

161 imply that the Te(1)-Tl distance at 3.49 Å in honeaite is likely to be a bond, albeit
162 contributing little bond valence to Te. Much longer Te-Tl distances in honeaite (> 3.9 Å) are
163 very unlikely to be bonds and if included would result in very unusual coordination for Te.
164 Thus, Tl can be considered to be octahedrally-coordinated. This coordination environment is
165 very different from that of Cs and Rb in CsAu_3S_2 , RbAu_3Se_2 and CsAu_3Se_2 , in which they are
166 octahedrally-coordinated by six Se (S) atoms.

167 Three gold telluride minerals related to honeaite are calaverite AuTe_2 , sylvanite
168 AuAgTe_4 and krennerite Au_3AgTe_8 (Dye and Smyth, 2012), all of which have general 1:2
169 stoichiometry (Au, Ag) Te_2 . Honeaite is the only gold telluride with essential thallium. While
170 calaverite, krennerite and honeaite have very different topologies, they share some features,
171 such as TeAu_3 pyramids and octahedrally-coordinated Au. However, krennerite has TeAu_3
172 pyramids that are much more distorted than those of honeaite. The 1:2 compounds have a
173 sheet of edge-sharing (Au,Ag) Te_6 octahedra; in sylvite and krennerite there are alternating
174 rows of AuTe_6 and AgTe_6 . In calaverite and sylvanite these sheets are planar with dative
175 bonding involving Te providing the only inter-sheet connectivity. In krennerite the sheets are
176 folded in a zigzag arrangement; again dative bonding involving Te is the only inter-sheet
177 connection. In contrast to these 1:2 sheet structures, honeaite does not appear to have dative
178 bonding; the only linkages between double-sheets are Tl-Au bonds.

179 It has been suggested that the short $\text{Te}\cdots\text{Te}$ distance of 2.84 Å in krennerite Au_3AgTe_8
180 may be associated with $\text{Te}\cdots\text{Te}$ inter-sheet bonding (Dye and Smyth, 2012). Indeed, no other
181 options appear to be possible for bonding between the sheets. Sylvanite AgAuTe_4 , another
182 sheet structure, has a $\text{Te}\cdots\text{Te}$ inter-sheet distance of 2.89 Å. Calaverite AuTe_2 , has a shortest
183 inter-sheet $\text{Te}\cdots\text{Te}$ distance of 3.20 Å. The shortest inter-sheet $\text{Te}\cdots\text{Te}$ and Te-Au distances in

honeaite are 4 Å, and as such are likely to be far too long to correspond to dative bonds.
Hence, the only significant inter-sheet bonding in honeaite is *via* the Tl-Au(2) bond (2.97 Å).

There are two synthetic silver-thallium-tellurides having an analogous Ag₃TlTe₂ stoichiometry: Ag₃TlTe₂ (Gelato et al. 1998) and Ag₁₈Tl₄Te₁₁ (Brun et al. 1992). Both have very different structures from each other and from that of honeaite. Ag₃TlTe₂ (orthorhombic, *Cmmm*) is an Ag-Te framework with strong compositional layering and Tl in [8]-coordination Tl(Ag₄Te₄). Ag₁₈Tl₄Te₁₁ (cubic, *F43m*) is an Ag-Te framework with channels occupied by Tl₆ octahedral clusters. In both compounds Te has high coordination ≥[6].

Finally, petzite Ag₃AuTe₂ (Haidinger 1845 cited by Frueh, 1959) and fischesserite Ag₃AuSe₂ (Bindi and Cipriani, 2004) have the same 3:1:2 stoichiometry but very different structures from that of honeaite.

Honeaite is a new gold telluride structure. The Tl site has four bonds to Au at ~ 3 Å and two long Tl-Te bonds. It may be possible for Au or Ag to occupy a similar, albeit more contracted, site. The possibility may, therefore, exist of a new class of 2:1 gold tellurides (Au, Ag)₂Te based upon the honeaite topology. We are not aware of any synthetic Au-Tl-Te phases, and so the wider significance of Tl for structural diversity in gold tellurides remains to be explored.

Acknowledgments

CJS acknowledges Natural Environment Research Council grant NE/M010848/1 Tellurium and Selenium Cycling and Supply. We thank Nigel Cook, Peter Leverett and an anonymous reviewer for their comments on this manuscript. Associate Editor David Hibbs is thanked for his efficient handling of the review process.

207

208 **References cited**

- 209 Brun, G., Moreau, J.M., Liautard, B., Allemand, J. (1992) A new phase in the Tl-Ag-Te
210 system: crystal structure of $\text{Tl}_4\text{Ag}_{18}\text{Te}_{11}$. Journal of Alloys and Compounds, 178, 85-
211 90.
- 212 Dye, M.D., Smyth, J.R. (2012) The crystal structure and genesis of krennerite, Au_3AgTe_8 .
213 Canadian Mineralogist, 50, 119-127.
- 214 Farrugia, L. J. (1999) WinGX: An integrated system of Windows Programs for the solution,
215 refinement and analysis of single-crystal X-ray diffraction data. Journal of Applied
216 Crystallography, 32, 837–838.
- 217 Frueh, A.J. (1959) The crystallography of petzite, Ag_3AuTe_2 . American Mineralogist, 44,
218 693-701.
- 219 Gelato, L.M., Penzo, M., Parthe, E., Cenzual, K. (1991) Inorganic structure types with
220 revised space groups I. Acta Crystallographica, B47, 433-439.
- 221 Klepp K.O., Weithaler C. (1996) The crystal structures of CsAu_3S_2 , RbAu_3Se_2 and CsAu_3Se_2
222 and their relationship to the CsCu_3S_2 structure type. Journal of Alloys and
223 Compounds, 243, 1-5.
- 224 Nechaev, S.V., Cook, N.J. (2000) A natural occurrence of Au_3TlTe_2 from the Maiskoe Au-
225 deposit, Ukraine. Neues Jahrbuch für Mineralogie, Monatshefte, Heft 12, 557–562.
- 226 Rice, C.M., Welch, M.D., Still, J.W., Criddle, A.J. and Stanley, C.J. (2015) Honeaite, IMA
227 2015-060. CNMNC Newsletter No. 27, 1229; Mineralogical Magazine, 79, 1229-
228 1236.

229 Rice, C.M., Welch, M.D., Still, J.W., Criddle, A.J. and Stanley, C.J. (2016) Honeaite, a new
 230 gold-thallium-telluride from the Eastern Goldfields, Yilgarn Craton, Western
 231 Australia. *European Journal of Mineralogy*, in press.

232 Schütte, W.J. De Boer, J.L. (1988) The incommensurately modulated structure of calaverite,
 233 AuTe_2 . *Acta Crystallographica*, B44, 486-494.

234 Sheldrick, G.M. (2008) A short history of SHELX. *Acta Crystallographica*, A64, 112-122.

235

236 Tunell, G. (1941) The atomic arrangement of sylvanite. *American Mineralogist*, 26, 457-
 237 477.

238 Wilson, A.J.C. (ed.) (1992) *International Tables for Crystallography, Volume C*. Kluwer
 239 Academic Publishers, Dordrecht.

240

241

242 **Figure Captions**

243 **Figure 1. (a)** The six-membered ring comprising fused chains of Te(1)Au_3 and Te(2)Au_3
 244 pyramids. **(b)** A pair of Te(1)Au_3 and Te(2)Au_3 pyramids with Te-Au bond lengths shown.
 245 The apex (Te) of the Te(1)Au_3 pyramid points up, that of the Te(2)Au_3 pyramid points down.
 246 **(c)** connected six-membered rings forming a corrugated sheet composed of alternating rows
 247 of Te(1)Au_3 and Te(2)Au_3 pyramids. **(d)** The same sheet as in (c) projected onto (001) and
 248 showing the nature of the sheet corrugation.

249

250 **Figure 2. (a)** Polyhedral representation of the honeaite structure showing corrugated layers
 251 composed of double-sheets of TeAu_3 pyramids with rows of thallium atoms (green spheres)

occupying grooves in sheet corrugations extending parallel to [001]. **(b)** A ball-and-spoke representation of (a) showing Tl-Au bonds (dashed black lines), Tl...Te distances (dashed red lines) and intra-sheet Au-Au bonds (solid blue lines).

Figure 3. (a) Coordination and bond distances ($\text{\AA}$) of Au and Tl atoms in honeaite. Au(1) and Au(3) form octahedra, whereas Au(2) is four-coordinate, including an inter-sheet Au-Tl bond (highlighted in red). Tl-Te distances are shown as dotted black lines and may not correspond to real bonds. Anisotropic displacement parameters are drawn at the 68% level. **(b)** Polyhedral connectivity of the three Au sites of honeaite. All atoms are drawn as spheres and central Au sites are numbered.

Figure 4. Fragment of the honeaite structure viewed slightly obliquely to the y axis and showing the very different polyhedral connectivities of Te(1)Au_3 (blue) and Te(2)Au_3 (grey) pyramids. Chains of TeAu_3 pyramids of the corrugated sheets extend parallel to [001]. For each pyramid the Te atom is located at the apex that is not bonded to other polyhedra. TlAu_4 groups are shown green and each shares two edges with Te(1)Au_3 pyramids, making a fourth longer inter-sheet bond to a Te(2)Au_3 pyramid of the adjacent double-sheet.

Figure 1

(a)

(b)

(c)

(d)

Figure 2

This is a 'preproof' accepted article for Mineralogical Magazine.
This version may be subject to change during the production process.
10.1180/minmag.2016.080.112

Table 1. Summary of the data collection and structure refinement of honeaite.

<i>Crystal data</i>	
Ideal chemical formula	Au ₃ TlTe ₂
Space group	<i>Pbcm</i>
<i>a</i> (Å)	8.9671(4)
<i>b</i> (Å)	8.8758(4)
<i>c</i> (Å)	7.8419(5)
<i>V</i> (Å ³)	624.14(6)
<i>Z</i>	4
<i>D_x</i> (Mg m ⁻³) calc	11.18
<i>μ</i> (mm ⁻¹)	105.05
<i>Data collection</i>	
Diffractometer	Xcalibur E (1K Eos detector)
Radiation, wavelength (Å)	MoK α , 0.71073
Crystal	Opaque rectangular plate
Max. Med. Min. dimensions (mm)	0.040 × 0.035 × 0.015
Temperature (K)	293(2)
Scan type, frame-width (°), frame-time (s)	ω , 1.0, 240
Absorption correction	Multi-scan
<i>T_{min}</i> , <i>T_{max}</i>	0.233, 1
Reflections used for cell, <i>I</i> > 7 σ (<i>I</i>)	1903
Reflections measured	11673
<i>R_G</i>	0.030
Independent reflections	1136
Independent reflections with <i>I</i> > 2 σ (<i>I</i>)	948
<i>R_{int}</i> (<i>mmm</i>)	0.058
$\theta_{\min}$, $\theta_{\max}$ (°)	3.22, 31.91
Index range	-12 ≤ <i>h</i> ≤ 13, -12 ≤ <i>k</i> ≤ 13, -11 ≤ <i>l</i> ≤ 11
Data completeness to 30° θ (%)	99.9
<i>Refinement</i>	
Reflections, restraints, parameters	1136, 0, 34
<i>R</i> ₁ [<i>I</i> > 2 σ (<i>I</i>)], <i>R</i> ₁ (all)	0.033, 0.046
<i>wR</i> ₂ [<i>I</i> > 2 σ (<i>I</i>)], <i>wR</i> ₂ (all)	0.050, 0.053
<i>GoF</i> (<i>F</i> ²)	1.087
Weighting scheme*	<i>a</i> = 0.0178
(Δ/σ) _{max}	<0.001
$\Delta\rho_{\max}$, $\Delta\rho_{\min}$ (e Å ⁻³)	2.1, -3.6

* SHELX (Sheldrick, 2008)

Table 2. Atom coordinates and displacement parameters U^{ij} ($\text{\AA}^2$) for honeaite.

atom	x/a	y/b	z/c	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}	U_{eq}
Tl	0.21609(5)	$\frac{3}{4}$	0	0.0196(3)	0.0263(3)	0.0255(3)	0.0043(2)	0	0	0.0238(1)
Au(1)	-0.13342(5)	$\frac{3}{4}$	0	0.0173(2)	0.0177(2)	0.0155(3)	0.0004(2)	0	0	0.0168(1)
Au(2)	0.54678(5)	$\frac{3}{4}$	0	0.0169(2)	0.0247(3)	0.0205(3)	-0.0064(2)	0	0	0.0207(1)
Au(3)	0.17523(5)	0.48884(5)	$\frac{1}{4}$	0.0156(2)	0.0161(2)	0.0186(3)	0	0	0.0018(2)	0.0167(1)
Te(1)	0.10918(9)	0.05618(8)	$\frac{1}{4}$	0.0143(4)	0.0152(4)	0.0150(4)	0	0	-0.0002(3)	0.0148(2)
Te(2)	0.53752(8)	0.55678(8)	$-\frac{1}{4}$	0.0150(4)	0.0145(4)	0.0154(5)	0	0	0.0003(3)	0.0150(2)

Table 3. Interatomic distances in honeaite (Å)

Au(1)		Tl	
Te(1) × 2	2.6173(5)	Au(2)	2.9653(7)
Au(2)	2.8677(7)	Au(3) × 2	3.0579(4)
Au(3) × 2	2.9117(3)	Au(1)	3.1341(7)
Tl	3.1341(7)	Te(1) × 2	3.4854(7)
Au(2)		Te(1)	
Te(2) × 2	2.6061(5)	Au(1) × 2	2.6173(5)
Au(1)	2.8677(7)	Au(3)	2.620(1)
Tl	2.9653(7)		
Au(3)		Te(2)	
Te(2)	2.6074(9)	Au(2) × 2	2.6061(5)
Te(1)	2.620(1)	Au(3)	2.6074(9)
Au(1) × 2	2.9117(3)		
Tl × 2	3.0579(4)		