

1 ***Tetracapsuloides bryosalmonae* abundance in river water**

2

3 Running page head: *T. bryosalmonae* abundance in river water

4

5 Fontes, I.^{1,2,3,4}, Hartikainen, H.^{3,4}, Holland, Jason W.², Secombes, Chris J.² and Okamura, B.^{1*}

6

7 ¹ Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK

8 ² Scottish Fish Immunology Research Centre, University of Aberdeen, Aberdeen AB24 2TZ, UK

9 ³ Eawag, Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland

10 ⁴ ETH Zürich, Institute of Integrative Biology (IBZ), Zürich, Switzerland

11

12 * Corresponding author: b.okamura@nhm.ac.uk

13

14 **Abstract**

15 *Tetracapsuloides bryosalmonae* is a myxozoan parasite of freshwater bryozoans and
16 salmonids, causing proliferative kidney disease in the latter. To date, detection of the parasite
17 requires collection of hosts and subsequent molecular or histological examination. The
18 release of infectious spores from both hosts offers an opportunity to detect the parasite in
19 water samples. We developed a SYBR[®] Green quantitative real-time PCR (qPCR) assay for
20 *T. bryosalmonae* in water samples which provides an estimation of bryozoan malacospore
21 numbers and tested the assay in three rivers in southern England over a period of five weeks.
22 The assay proved to be both highly sensitive and specific to the parasite, detecting low levels
23 of spores throughout the study period. Larger volume samples afforded greater detection
24 likelihood, but did not increase the number of spores detected, possibly as a result of low and
25 patchy spore distributions and lack of within-site replication of large volume samples. Based
26 on point-measurements, temperature was positively associated with the likelihood of
27 detecting spores, possibly reflecting the temperature dependence of spore shedding from
28 bryozoan hosts. The presence of *T. bryosalmonae* in water samples was predominantly
29 influenced by spatial (sites within rivers, amongst rivers) and temporal (sampling dates)
30 factors, while the latter also influenced C_q values and spore abundance. Environmental
31 monitoring for infectious stages can complement traditional methods, providing faster and
32 easier detection and avoiding potentially prolonged searching, collecting and destructive
33 sampling of invertebrate and vertebrate hosts.

34

35 **Keywords:** Proliferative Kidney Disease; Myxozoa; qPCR; environmental DNA; disease risk

36 **Introduction**

37

38 Emerging aquatic diseases pose threats to biodiversity, conservation and sustainable use of
39 freshwater resources (Okamura & Feist 2011). Monitoring parasites and pathogens over
40 appropriate temporal and spatial scales is therefore crucial for understanding and predicting
41 the conditions that lead to disease outbreaks. However, detecting infections in the absence of
42 clinical disease and mortality can be challenging and problematic. For example, for many fish
43 diseases, detection involves destructive sampling of already threatened host species and may
44 require numerous individuals to be killed to gain confidence in the results. Time-consuming
45 histopathology or tissue-targeted molecular approaches may then be required to verify
46 parasite presence. As an alternative approach and a complementary tool, molecular detection
47 of parasite DNA in environmental samples is increasingly employed in marine and freshwater
48 environments (Audemard et al. 2006, Hung & Remais 2008, Strand et al. 2014). Thus, the
49 detection and quantification of disease agents in environmental samples offers a unique
50 potential to inform on the ecology and epidemiology of host-parasite interactions by
51 circumventing traditional parasitological approaches. With these advantages in mind, we
52 have developed a quantitative real-time PCR (qPCR) assay to detect and characterise, from
53 water samples, the abundance of *Tetracapsuloides bryosalmonae*. This myxozoan causes
54 proliferative kidney disease (PKD) in salmonids – a disease that has been increasing in
55 prevalence and severity, particularly in fish farms, and whose distribution has been expanding
56 with environmental change.

57

58 *Tetracapsuloides bryosalmonae* is an endoparasite of freshwater bryozoans and salmonids,
59 causing PKD in both wild and farmed fish in Europe (Wahli et al. 2007) and North America
60 (Ferguson & Needham 1978). *Tetracapsuloides*, *Buddenbrockia* and several undescribed
61 species form the Malacosporea (Fiala et al. 2015), a small and early diverging clade of
62 myxozoans (Canning et al. 2000). *T. bryosalmonae* spores (referred to as malacospores: Feist
63 et al. 2015) released in the urine of fish measure some 16µm in width and 14 µm in height
64 (Kent & Hedrick 1986, Hedrick et al. 2004, Bettge et al. 2009) and are infective to freshwater
65 bryozoans (Morris & Adams 2006, Grabner & El-Matbouli 2008). In bryozoans the parasite
66 forms sacs (up to 350µm in diameter) filled with many thousands of spherical spores of
67 approximately 20µm in diameter (Canning et al. 2000, McGurk et al. 2005, Okamura et al.
68 2011). Malacospores released from sacs are ejected from bryozoans and remain infectious to

69 fish for 12-24h (Feist et al. 2001, De Kinkelin et al. 2002). The smaller spores of *T.*
70 *bryosalmonae* released from fish possess two capsulogenic cells, a single sporoplasm and two
71 valve cells (Morris & Adams 2008). The larger spores released from bryozoans have four
72 capsulogenic cells, two sporoplasms, and some eight or ten valve cells (Feist et al. 2015).
73 Little is known about the timing of malacospore release from bryozoan and fish hosts nor of
74 variation in spore abundance in natural systems. In bryozoan hosts, spore production has been
75 observed to occur predominantly in spring and autumn (Tops et al. 2006), which should lead
76 to increased spore concentrations in water at these times. However, because naïve fish
77 become infected in other seasons, spores released from bryozoans are likely to be present
78 year-round (Gay et al. 2001). The presence of sporogonic stages (pseudoplasmodia in kidney
79 tubules) in naturally-infected fish has been described as rare and spores are estimated to be
80 released in numbers that will be greatly diluted in the natural environment (maximum
81 concentration estimate = 120 spores/ml urine) (Hedrick et al. 2004).

82

83 Detection of *T. bryosalmonae* currently includes searching for and collection of patchily
84 distributed freshwater bryozoans, followed by qualitative PCR to confirm infection or
85 examining dissected bryozoans for spore-producing sacs. Detection of infection in wild fish
86 typically involves electrofishing, dissection and subsequent histopathology or conducting
87 PCR/qPCR of fish tissues (e.g. Grabner & El-Matbouli 2009, Kumar et al. 2013). qPCR
88 assays to detect and quantify other myxozoans in water samples have been developed for
89 *Ceratonova shasta* (Hallett & Bartholomew 2006), *Parvicapsula minibicornis* (Foott et al.
90 2007), *Henneguya ictaluri* (Griffin et al. 2009), *Ceratonova puntazzi* (Alama-Bermejo et al.
91 2013) and *Kudoa yasunagai* (Ishimaru et al. 2014) (for review, see Fontes et al. 2015). The
92 development of qPCR to enable molecular detection and quantification specific to *T.*
93 *bryosalmonae* spores in water samples would help to avoid or reduce the labour-intensive
94 approaches currently employed to ascertain the presence and abundance of *T. bryosalmonae*
95 in water bodies.

96

97 The aims of this study were to: (1) Develop a novel *T. bryosalmonae* SYBR[®] Green qPCR
98 assay; (2) Use the assay to quantify *T. bryosalmonae* spores in rivers with known PKD
99 occurrence in southern England over time; (3) Determine how the detection and abundance of
100 *T. bryosalmonae* spores are affected by sample location, sample volumes, time of sampling,
101 and point-measurements of water temperature and flow.

102

103 **Materials and Methods**

104

105 *Sampling sites*

106 The sites selected for this study are on the Rivers Avon and Itchen in Hampshire and the
107 River Dun in Berkshire. The rivers represent spring-fed chalk stream systems that harbour
108 wild brown trout (*Salmo trutta*) populations in lowland habitats in southern England. The
109 rivers also provision rainbow trout (*Oncorhynchus mykiss*) fish farms that sustain regular
110 PKD outbreaks. The bryozoan, *Fredericella sultana* occurs abundantly in the rivers growing
111 as dense stands of colonies attached to submerged roots of riparian alder and willow trees.
112 Water was sampled for qPCR studies near three separate *F. sultana* populations that were
113 known to be infected (Fontes 2015) (on three different tree root systems) in each river.

114

115 *Water sampling*

116 Water samples were collected every Monday for five weeks in the three rivers during the
117 period when high numbers of spores were expected to be released from infected bryozoans
118 (May-June) (Tops 2004). Sampling of the Rivers Avon and Dun commenced on 14/05/12 and
119 of the River Itchen on 13/05/13. Water samples were collected from approximately 30 cm
120 below the water surface and 1 m downstream from the tree root systems (see Figure 1). One
121 2L plastic bottle was filled downstream from each root system by submerging the bottle to
122 collect incoming flow. The submerged bottle was oriented upstream and slowly moved from
123 side to side to collect water across approximately a 1.5 m stretch of the river, perpendicular to
124 the river bank. In addition to these 2L samples, a bucket was used to fill one 24L plastic
125 container with water collected 1m downstream from the most downstream of the three roots
126 in each river (Figure 1). Hence, for each sampling point a total of four water samples were
127 taken in each river: three 2L samples at each root and one 24L sample at the most
128 downstream root. At each sampling date, point-measurements of water temperature and water
129 flow (mean velocity over 60 s, using an electromagnetic open channel flow meter (Model
130 8008/801, Valeport Ltd., Totnes, UK) were noted at each root. The samples were stored at
131 4°C in the original collection containers and filtered within 24 h of collection.

132

133 *Water filtration and DNA extraction*

134 The 24L samples were pre-filtered through a 30 µm mesh in the field as the containers were
135 filled. All samples were then filtered in the laboratory onto cellulose nitrate filter membranes
136 (3 µm pore size, 142 mm diameter; Sartorius Stedim Biotech GmbH, Goettingen, Germany)
137 at 1 bar using a pressure filtration system (Sartorius Stedim Biotech GmbH, Goettingen,
138 Germany). Filter papers were scraped with a razor blade and the scrapings placed in a 1.5 ml
139 micro-centrifuge tube and stored at -80°C. The filtration system was rinsed thoroughly with
140 deionised water and razor blades were sterilised using EtOH before processing each sample.
141 Samples were freeze-dried at -56°C to remove excess water and DNA was extracted using an
142 UltraClean® Soil DNA kit (MO BIO Laboratories Inc., Carlsbad, California, USA). The 50
143 µL eluted DNA was then preserved at -20°C and defrosted temporarily for screens.

144

145 ***qPCR standards***

146 A 244 bp fragment of the *T. bryosalmonae* small subunit rDNA (SSU rDNA) gene was
147 amplified from a genomic DNA sample derived from 85 mature spores obtained from sacs
148 dissected from field collected colonies (from the River Avon) using the specific primers,
149 514F_new (5'-ATTCAGGTCCATTCGTGAGTAACAAGC-3') (Hartikainen et al. 2013) and
150 776R (5'-GCTGATACACCCAATTAAGGGCAG-3') (Morris et al. 2002). The resulting
151 PCR product was purified and concentration measured using a Thermo Scientific NanoDrop
152 8000 Spectrophotometer (in ng/µl), adjusting it to 1 nM in 1 ml of TE buffer based on the
153 mean molecular weight of a base pair (i.e. 660 Da). A 1:10 serial dilution of the standardised
154 1 nM solution was performed and used as a template for the qPCR standard curve (n = 7
155 concentrations) with the primers described below (518F_Q and 680R_Q nest completely
156 within 514F_new and 776R). The seven standards used encompassed the full range of
157 samples tested. All quantification cycle (C_q) values were determined using a fixed threshold
158 normalised fluorescence of 0.1 (obtained manually) across all runs. The standard curve was
159 applied to all runs using the first standard (1e-12 mol/L) to normalise each respective run.

160

161 ***qPCR assay***

162 To detect and quantify *T. bryosalmonae* SSU rDNA, a SYBR® Green qPCR assay was
163 developed, using species specific primers 518F_Q (5'- CAGGTCCATTCGTGAGTAACAA-
164 3') and 680R_Q (5'- TGCCTCCTTAGTTAGGTAGACAAA-3' (Sigma-Aldrich®, Poole,
165 UK; primers were purified using the desalted method) and targeting a 182 bp fragment of the
166 *T. bryosalmonae* SSU rRNA gene. Primers were designed based on inspection of

167 comprehensive alignment of all known malacosporean 18S SSU rDNA sequences. Dimer
168 formation and primer quality were checked using NetPrimer
169 (<http://www.premierbiosoft.com/netprimer/>) and via blast searches against the NCBI
170 GenBank database (<http://www.ncbi.nlm.nih.gov/genbank/>). Primers developed for the assay
171 target a portion of the SSU rRNA gene, which is present in the parasite genome as a tandem
172 repeat unit. Focussing on multi-copy genes provides an advantage over single-copy genes in
173 terms of detection sensitivities. Furthermore, as the number of tandem copies is
174 approximately the same in each cell, SSU rDNA avoids potential biases associated with
175 mitochondrial targets, which, although present in multiple copies, may vary in number
176 depending on developmental stage, cell type and physiological state. The final volume of the
177 qPCR mix was 10 μ L that was comprised of: 1 μ L of template DNA, 1x of Rotor-Gene
178 SYBR[®] Green PCR Master Mix (QIAGEN GmbH, Hilden, Germany), 1 μ M of each primer
179 and molecular grade water (Fisher Scientific). The relative concentration of each primer was
180 optimised in a test of nine combinations from 0.05 - 1 μ L of either forward or reverse primer.
181 The optimal combination was 1 μ L of each primer as this was the one that produced the
182 lowest C_q value and the highest fluorescence (see Online Table 1). A CAS-1200[™] pipetting
183 robot (Corbett Life Science, Mortlake, Australia) was used to prepare and dispense the master
184 mix and template DNA into 0.2 ml clear PCR tube strips (QIAGEN GmbH, Hilden,
185 Germany). Each qPCR run included the following (in duplicate): (1) Negative control (water
186 only); (2) Positive control (gDNA from a single *T. bryosalmonae* spore released from a
187 bryozoan host); (3) Water samples to be tested; and (4) Seven qPCR standards (1e-12 to 1e-
188 18 mol/L). A sample would only be considered to be positive if both duplicate reactions were
189 positive. Reactions were performed in a Rotor-Gene[™] 6000 real-time PCR machine (Corbett
190 Life Science, Mortlake, Australia) and runs analysed using the Rotor-Gene[™] 6000 Series
191 Software 1.7 (Corbett Life Science, Mortlake, Australia). The thermal cycling conditions
192 were: 95°C for 5 min, followed by 45 cycles of: denaturation at 95°C for 5 s; and
193 annealing/extension at 60°C for 10 s (as recommended by QIAGEN for SYBR green assays).
194 Data were acquired at the end of each cycle on the green dye channel (470 \pm 10 nm
195 excitation, 510 \pm 5 nm detection, 9.67 gain). A melting curve between 74-95°C was run at the
196 end of each qPCR run. Water samples without a fluorescent signal were re-tested alongside
197 negative and positive control samples. Intra-assay variability (repeatability) was calculated as
198 the coefficient of variation (CV) for concentration variance (standard deviation [SD]) of
199 seven standards and all eDNA samples. Inter-assay variability (reproducibility) was

200 calculated as the CV of concentrations of each of the seven standards between four runs. All
201 runs were performed by the same operator.

202

203 ***qPCR assay sensitivity and specificity***

204 To test the sensitivity of the assay, serial dilutions (1:10 – 1:100,000) of two positive and
205 three negative (undiluted; used as controls) river water samples were analysed. We
206 considered the limit of detection (LOD) of the assay to be at the highest C_q value after which
207 there was no fluorescence detected in dilutions of positive samples (Francois et al. 2003,
208 Hallett & Bartholomew 2006). This definition of LOD is conservative for detection, as it
209 minimises the chances of false positives.

210

211 To test that the primers used were specific for *T. bryosalmonae*, we undertook qPCR analysis
212 using the following range of templates: other malacosporean samples (*Buddenbrockia*
213 *allmani*; *B. plumatellae*; *Buddenbrockia* species 2 and novel lineages 1-3 (Hartikainen et al.
214 2014); *T. bryosalmonae* sacs and respective spores; *Fredericella sultana* colonies not infected
215 by *T. bryosalmonae*; *F. sultana* colonies with covert and overt *T. bryosalmonae* infections;
216 uninfected and *T. bryosalmonae*-infected rainbow trout kidney tissue, the latter showing
217 clinical signs of PKD. qPCR products of samples exhibiting a fluorescent signal were verified
218 by direct sequencing on an ABI PRISM[®] 3700xl DNA analyser (Applied Biosystems[™],
219 Foster City, USA) using BigDye v1.1 chemistry.

220

221 ***Inhibition testing***

222 To assess the presence of PCR inhibitors, qPCR amplification of an internal positive control
223 (IPC) was compared in reactions containing eDNA extract to those only containing DNA-free
224 water (Sigma-Aldrich[®], Poole, UK). This test was carried out for a subset of samples (Online
225 Table 2). A total of seven river water samples (3 x 24L and 4 x 2L) were randomly selected
226 for this test. A synthetic IPC template was designed (5'-
227 GTATTCCTGGTTCTGTAGGTTGAGCGTAAAACGACGGCCAGTGAATTGTAATACG
228 ACATGGTCATAGCTGTTTCCCGATACGGAAGTCCAGTCACAT -3') (Microsynth AG;
229 97 bp, purified using the desalted method), including two priming sites with no known
230 homology to published sequence data. The IPC template concentration was adjusted to 1 nM
231 (using Qubit[®] 2.0 Fluorometer) and stored in TE buffer. A serial dilution (1:10) of the
232 standardised solution was performed and the standard 1e-14 mol/L used as the IPC in a qPCR

233 assay with primers MIMf (5' - GTATTCCTGGTTCTGTAGGTTGAGC -3') and MIMr (5'
234 ATGTGACTGGACTTCCGTATCG -3'). A QIAgility pipetting robot (QIAGEN GmbH,
235 Hilden, Germany) was used to prepare and dispense the master mix and template DNA, as
236 well as the eDNA sample potentially containing the inhibitors. Each reaction containing IPC
237 and eDNA sample was run in duplicate. In control reactions the eDNA was replaced with
238 DNA-free water and ran as six replicates. Total reaction volume was 10 μ L containing: 1e-15
239 mol/L of IPC, 1 μ L of a river water sample or DNA-free water, 1x SYBR® Select Master
240 Mix (Applied Biosystems™, Foster City, USA) and 0.4 μ M of each primer (MIMf and
241 MIMr). The thermal cycling conditions were as follows: 50°C for 2 min; 95°C for 2 min; and
242 40 cycles of denaturation at 95°C for 15 s and annealing/extension at 60°C for 1 min. This test
243 was run on a 7500 Fast Real-Time PCR System (Applied Biosystems™, Foster City, USA)
244 using a standard ramp speed and analysed using the 7500 Software version 2.0.6 (Applied
245 Biosystems™, Foster City, USA). C_q values were determined using a fluorescence threshold
246 of 0.1 (obtained manually). Significant inhibition in a sample was defined as a difference of >
247 three cycles between mean C_q values of IPC reactions with and without eDNA (Hartman et
248 al. 2005). In such cases, samples were not used for further analyses as results of the *T.*
249 *bryosalmonae* assay may appear biased due to inhibition. In addition to this test, the serial
250 dilutions of the river water samples used in the sensitivity test were run with and without the
251 inclusion of bovine serum albumin (BSA; 250 ng/ μ L final concentration) in the master mix.

252

253 ***Estimating rDNA content of malacospores from bryozoans***

254 To estimate the number of *T. bryosalmonae* spores present in river water samples, we used
255 forceps and a needle to release bryozoan malacospores from a sac. Spores were then rinsed in
256 deionised water and individually pipetted, using a micro-injector (at 100-400 x
257 magnification), into 1.5 ml micro-centrifuge tubes and stored at -80°C. DNA from each spore
258 was extracted using a DNeasy® Blood & Tissue kit (QIAGEN GmbH, Hilden, Germany),
259 eluted in 200 μ L TE buffer, and quantified using the qPCR protocol described above for
260 water samples. Six individually extracted spores were used to estimate the SSU rDNA
261 content of a single mature malacospore obtained from a bryozoan host. No malacospores
262 from fish were available.

263

264 ***Statistical analysis***

265 All statistical analyses were performed using R (version 2.15.1) (R Core Team 2014). Welch
266 two sample t-tests were used to test differences in C_q values assessed in: 1) IPC reactions
267 spiked with and without river water; and 2) IPC reactions spiked with 2L and 24L river water
268 samples. Parasite presence and abundance in river water samples was related to potential
269 explanatory variables (sampling type (2L or 24L); point-measurements of water temperature;
270 and water flow) using Generalised Linear Mixed Models (GLMMs) following the methods
271 described in Zuur et al. (2009). The lme4 package (version 1.1-7) was used to analyse the
272 parasite presence/absence data, assuming a binomial error distribution (Bates et al. 2013). For
273 parasite abundance, the nlme package (version 3.1-117 was used, assuming a Gaussian error
274 distribution (Pinheiro et al. 2014). Random effects models with no fixed factors were used to
275 determine the optimal random effects structure using restricted maximum likelihood
276 estimation (REML, parasite presence) or maximum likelihood estimation (ML, parasite
277 abundance). Univariate analyses were then performed on each explanatory variable and those
278 with P-values below 0.25 were included in a maximal model using ML following a visual
279 check to remove any covariates that were strongly correlated. Non-significant variables and
280 interactions were eliminated in a stepwise fashion, removing the least significant
281 relationships first until only variables significant at $P < 0.05$ remained. Random intercept and
282 slope models with intercept values of the significant fixed effects were evaluated and only
283 retained if they led to a significant reduction in a model's log-likelihood.

284

285 **Results**

286

287 *qPCR sensitivity and specificity testing*

288 The linear standard curve (Online Figure 1) had a slope of -3.37, a correlation coefficient (R^2)
289 of 0.998 and an amplification efficiency of 98%. Although this standard curve was applied to
290 all the runs using the first standard as a reference, the standards curves included in each run
291 performed well with efficiency being higher than 96% and an R^2 ranging from 0.989 - 0.999.
292 The dilution series of two known positive samples (both replicates fluorescing) indicated that
293 the fluorescence signal was lost at an approximate mean C_q value of 31.01 and 27.46 in each
294 qPCR-positive water sample (Figure 2 and Online Table 3). This suggests that the LOD for
295 this assay is 31 C_q as this was the highest value obtained for the two positive samples. The
296 mean concentration of parasite rDNA in a bryozoan malacospore, based on six individual
297 spores, was $1.96e-18$ mol/L ($\pm 1.69e-19$ SD) (i.e. 25.83 C_q) in 200 μ L, which equates roughly

298 to 0.005 spores per qPCR reaction based on a reaction volume of 10 μ L. A value of 31 C_q
299 equates to 0.089 spores in 50 μ L of eluted DNA ($1.75e-19$ mol/L - calculated using the
300 standard curve equation presented in Online Figure 1 [$C_q = 10^{(-0.297 * 31 - 9.551)}$]) or
301 0.0018 spores per qPCR reaction (spore numbers calculated based on the concentration of 1
302 spore [$1.96e-18$ mol/L]). Samples that were negative when undiluted did not exhibit any
303 fluorescence at any dilution (see Figure 2).

304
305 Melting curves produced two peaks. The first was at $\sim 77.3^\circ\text{C}$, corresponding to the presence
306 of primer dimer. The second peak, at $\sim 84.5^\circ\text{C}$, corresponded to amplification of the target
307 template. Negative samples and controls produced the first peak, in positive samples only the
308 second peak was present. Although an LOD of 31 C_q was implied by the sensitivity analysis,
309 samples with a sub-LOD concentration were found in 22% of the river samples ($n = 60$), with
310 C_q values up to 34.5 (corresponding to 0.015 spores in 50 μ L of eluted DNA ($2.96e-20$
311 mol/L) or 0.0003 spores per qPCR reaction). Differences between the sensitivity analysis and
312 the tested samples suggest that a C_q of 31 is the limit of quantification (LOQ) which, for
313 complex samples, is generally 5 - 10 times higher than the absolute LOD (Berdal & Holst-
314 Jensen 2001). The assay exhibited both low to high repeatability and high reproducibility.
315 The former is supported by substantial variation in intra-assay variance for eDNA samples
316 (CV range = 3.21 – 74.70%) and a low intra-assay variance for standards (CV range = 0.08 –
317 39.24%). The latter is supported by the small inter-assay variance for the concentrations of all
318 seven standards (CV range = 5.28 – 10.03%; see Online Table 4). Therefore, we set the LOQ
319 in the qPCR assay at 31 C_q , which is six times the concentration of the absolute LOD (34.5
320 C_q). For statistical analyses, the LOQ was used as the cut-off for parasite presence and
321 abundance.

322
323 The qPCR assay was highly specific to the presence of *T. bryosalmonae*. No amplification
324 was observed in negative samples (i.e. uninfected *F. sultana* colony, uninfected rainbow trout
325 kidney and *Buddenbrockia* samples) and all amplified products were verified by sequence
326 analysis as belonging to *T. bryosalmonae*. The C_q results for *T. bryosalmonae* were as
327 follows: one bryozoan malacospore (mean = 27.45, SD = 0.08, $n = 6$); and sacs with an
328 unknown number of spores at potentially different developmental stages (mean = 10.43, SD =
329 0.37). Infected host material produced the following C_q results: overtly infected colony

330 (mean = 15.34, SD = 0.31); covertly infected colony (mean = 24.13, SD = 0.11); and PKD-
331 affected rainbow trout kidney (mean = 18.35, SD = 0.00).

332

333 ***Inhibition testing***

334 A significant increase in C_q values during IPC amplification was observed when river water
335 was added to reactions (Welch Two Sample t-test: $t = 2.942$, d.f. = 8.595, $P = 0.017$),
336 indicating the presence of PCR inhibition. Spiking of an IPC reaction with river water
337 samples (Online Table 2) increased C_q values on average by 1.112 (C_q with river water,
338 mean = 17.452, SD = 0.341, $n = 7$; C_q without river water: mean = 16.340, SD = 0.166, $n =$
339 6). As differences were lower than three cycles all samples were used in the subsequent
340 analyses. Apparent effects of inhibition were larger in 2L than 24L samples, with
341 significantly higher C_q values in 2L samples (mean = 18.005, SD = 0.814, $n = 4$) than in 24L
342 samples (mean = 16.715, SD = 0.152, $n = 3$) (Welch Two Sample t-test: $t = 3.102$, d.f. =
343 3.279, $P = 0.047$). The standard deviation between replicates of all IPC reactions was small
344 (mean = 0.219, SD = 0.188, $n = 8$) and the dissociation melting curves of positive reactions
345 produced a sharp peak at $\sim 79^\circ\text{C}$. The addition of BSA to PCR reactions did not improve
346 amplification success (see Online Table 3).

347

348 ***Parasite presence and abundance in river water***

349 A total of 60 water samples (20 for each river) were collected (15 x 2L samples and 5 x 24L
350 samples per river). The mean temperature of five point measurements was not significantly
351 different between the Rivers Dun and Itchen, although it differed significantly between the
352 Rivers Avon and Itchen and between the Rivers Avon and Dun (see Table 1). Hence, the
353 River Avon was the warmest of the three rivers during the sampling period. Point-
354 measurements of water temperatures in the River Itchen remained lower and more stable
355 ($10.5 - 12.6^\circ\text{C}$) throughout the study than in the other two rivers (Avon: $12.0 - 18.4^\circ\text{C}$, Dun:
356 $10.5 - 15.9^\circ\text{C}$), where temperatures peaked in the third week (Figure 3a-c). Temperature did
357 not vary significantly between roots within a river, although water flow did (see Table 1),
358 particularly in the River Avon (see Figure 3a-c). There was no clear link between water flow
359 and temperature, the exception being in the River Avon when both environmental variables
360 peaked in the third week.

361

362 Almost half of the samples tested positive for *T. bryosalmonae* (40%). *T. bryosalmonae* was
363 detected in water samples taken each week in the Rivers Avon and Dun, but was only
364 detected in the River Itchen samples collected during the last two weeks of sampling (Figure
365 4). Parasite DNA was rarely detected in all samples from a given river on a given date. The
366 overall proportion of positive samples was higher in the River Avon (60%), than in the Rivers
367 Dun (40%) and Itchen (20%). The presence of *T. bryosalmonae* in water samples was
368 predominantly influenced by the individual rivers, root systems, and sampling dates. These
369 factors were subsequently included as random effects in mixed models to assess the
370 significance of fixed explanatory variables (sample volume, flow and point-measurements of
371 water temperature). Parasite DNA was detected more often using the 24L than 2L sampling
372 method. The 2L samples were 0.11 times (odds ratio) less likely to contain *T. bryosalmonae*
373 than 24L samples acquired on the same date. A total of 29% and 73% of 2L and 24L samples,
374 respectively, were positive. Temperature also had an effect on the presence of *T.*
375 *bryosalmonae* in the river water, with a unit increase in temperature increasing the likelihood
376 of presence by 1.60 times (odds ratio) (see Table 1).

377

378 C_q values in the river water samples were very close to the LOD, particularly those of 2L
379 samples (mean = 29.498, SD = 1.332, n = 13). Although not significant, C_q values decreased
380 and variation increased slightly in the 24L samples (mean = 28.989, SD = 1.355, n = 11).
381 Sampling date explained most of the variation in C_q values and none of the explanatory
382 variables were significantly associated with C_q values (see Table 1).

383

384 The estimated numbers of spores (converted from the template concentrations) ranged from
385 0.15-3.56 in the 2L sampling method (mean = 0.623, SD = 0.912, n = 13) and from 0.19-4.46
386 in the 24L sampling method (mean = 0.894, SD = 1.214, n = 11) (Figure 4 and Online Table
387 5 for the corresponding C_q values). Both methods detected up to 4 spores per water sample
388 but most positive samples using both methods contained less than one spore. In the River
389 Avon, *T. bryosalmonae* was more likely to be present in water sampled near the root system
390 furthest downstream, but no detectable pattern in spore detection relative to root systems was
391 found in the Rivers Dun and the Itchen (Online Table 5). The lower detection frequency in
392 the River Itchen was notable and was possibly associated with lower temperatures in this
393 river. Although not significant (see Table 1), spore numbers and variation in spore numbers

394 increased slightly in the 24L samples relative to the 2L samples. Sampling date explained
395 most of the variation in spore number.

396

397 **Discussion**

398

399 ***A new qPCR assay for T. bryosalmonae***

400 The novel *T. bryosalmonae* SYBR[®] Green qPCR assay to detect and quantify spores in water
401 samples performed consistently in three river systems over space and time. The LOD of the
402 assay was 34.5 C_q and the LOQ was 31 C_q, corresponding to 0.0003 and 0.0018 spores per
403 qPCR reaction, respectively. The assay was both sensitive and reliable, quantifying the
404 estimated SSU rDNA content of 0.005 spores consistently in six biological replicates (mean
405 = 25.83, SD = 1.24). The number of spores detected by the 2L sampling method ranged from
406 approximately 0.05 to 3.56 and by the 24L method from 0.02 to 4.46 spores. We suggest that
407 reasons why the larger samples did not detect more spores include patchiness of spores and
408 lack of replication within sites of large volume samples. We provide below further discussion
409 on how inhibition and spore quantification methods may affect our estimates.

410

411 ***Inhibition testing***

412 Detection of pathogens in natural water samples can be severely limited by PCR inhibition
413 due to substances such as calcium and humic acids (Opel et al. 2010). We developed an IPC
414 molecule and associated primers which can be used to test for the presence of inhibition in
415 reactions run alongside the quantification reactions. Although the C_q values increased
416 significantly when river water was added, the effect size was deemed to be low (1.112
417 cycles). Hartman et al. (2005) only consider a shift in C_q values of \geq three cycles as a sign of
418 inhibition. In our case no 24L samples showed significant inhibition, whereas 2L samples
419 from the Rivers Dun and Avon (but not Itchen) showed signals of low level inhibition. This
420 result is unexpected as large volume samples would be expected to suffer more from
421 accumulation of inhibitory compounds. In this study, the processing of the 2L and 24L
422 samples differed by pre-filtration through a 30 μ m mesh, which was only applied to the 24L
423 samples. It remains untested whether this may have removed particles carrying inhibitor
424 compounds and could explain the lower inhibition signal in the 24L samples. It should be
425 noted that these results are based on a low number of samples and thus interpreting patterns
426 between sample volumes is difficult. Nevertheless, the results underlie the conclusion that the

427 effects of inhibition in our samples are likely negligible. Also, no improvement to the final
428 workflow was seen following the dilution of river water samples, nor by the addition of BSA,
429 suggesting that the inhibition present in our river water samples is minimal as shifts of one
430 cycle can occur between runs and instruments. We recommend that an IPC is always
431 included to monitor the presence of inhibition in eDNA samples and may even be
432 multiplexed into sample assays on platforms allowing fluorescence detection on multiple
433 channels.

434

435 ***Detecting and quantifying T. bryosalmonae spores in river water***

436 *T. bryosalmonae* was detected in at least one water sample on all sampling dates in the Rivers
437 Dun and Avon but was not detected in the River Itchen until the fourth sampling date.

438 Estimated spore numbers were consistently low. The presence of *T. bryosalmonae* in water
439 samples was predominantly influenced by individual rivers, roots within rivers, and sampling
440 dates. However, despite the potential confounding effects of such spatial and temporal
441 variation, we were able to gain some insights into factors that may influence spore presence
442 and abundance. For example, we found that sampling date explained most of the variation in
443 spore number with few or no spores being detected in the early sampling periods in both 2L
444 and 24L samples from each river. In addition, we found that *T. bryosalmonae* was 1.6 times
445 more likely to be detected by qPCR given a unit increase in temperature - a result in keeping
446 with temperature-induced development and release of spores from bryozoans (see Tops &
447 Okamura 2003, Tops et al. 2006). However, it should be noted that the water temperature
448 measurements were only taken once a week rather than continuously. Although water flow
449 was found to be highly variable amongst rivers and roots, we found no effect of flow on the
450 detection or quantification of parasite spores. Foott et al. (2007) similarly found water flow to
451 have no influence on the detection of spores of the myxozoans *C. shasta* and *P. minibicornis*
452 in the Klamath River in California

453

454 It should be noted that malacospores released from bryozoans and fish differ in the number of
455 constituent cells. Furthermore, it is likely that these cells vary in ploidy levels. On the basis of
456 cell number, the rDNA content of fish malacospores may be estimated as $\leq 50\%$ than that of
457 bryozoan malacospores. However, fish malacospores are diploid while at least a proportion or
458 perhaps all of the cells in bryozoan malacospores are haploid (see Canning et al. 2007 for
459 discussion of ploidy of cells comprising bryozoan malacospores). Until both ploidy levels of

460 cells and cell numbers of malacospores are fully understood, our approach provides a direct
461 estimate of bryozoan malacospore concentrations but an underestimate of fish malacospore
462 concentrations. Ignoring unrelated copy number variation between spore states and cells and
463 applying the most conservative scenario based on known and proposed states (i.e. diploidy of
464 all fish malacospore cells and no secondary cell within the sporoplasm [Morris & Adams
465 2008] and diploidy of all bryozoan malacospore cells apart from haploid sporoplasms which,
466 however, do contain an internal haploid secondary cell [Canning et al. 2007]), the rDNA
467 content of fish malacospores would be some 46.7% less than those of bryozoan
468 malacospores. Thus fish malacospore rDNA concentrations would be underestimated by
469 approximately 100%.

470

471 A consideration of the dominant spore type in our samples is important given the above
472 variation in genomic DNA content of spores. Although timing of spore release from fish
473 hosts is unknown, spores are released in low numbers in fish urine (Hedrick et al. 2004) and
474 will be greatly diluted. It is therefore unlikely that spores deriving from fish substantially
475 contributed to the patterns detected here, particularly as we collected water directly
476 downstream from bryozoan populations known to sustain infections of *T. bryosalmonae*.

477

478 The 24L sampling method was more efficient than the 2L sampling method at detecting *T.*
479 *bryosalmonae* with detection more likely for the larger volume samples taken on the same
480 dates and at the same root systems. However, the numbers of spores estimated by qPCR for
481 2L and 24L samples were comparable (from 0.15 - 4.5). Detection of parasite DNA was
482 expected on all sampling dates based on our knowledge of spore development in bryozoans in
483 the field sites under investigation and results of previous studies quantifying myxozoans in
484 2L water samples by qPCR (Hallett & Bartholomew 2006). This was generally the case for
485 samples from the Rivers Dun and Avon, although not always for all samples on each date.
486 Lack of detection in some replicate samples is likely to represent false negatives due to
487 failure to detect or to capture spores in the samples. Since the assay was shown to be highly
488 sensitive and inhibition was low, the latter seems the most likely explanation - a premise
489 supported by the generally low spore concentrations in each river. The lower incidence of
490 false negatives in the two rivers with the 24L sampling technique (10%; n = 10) when
491 compared to the 2L method (63%; n = 30) provides additional support for this inference.
492 However, the consistent lack of detection of *T. bryosalmonae* in qPCR assays on River Itchen

493 water collected during the first three weeks suggests true absence during this period, possibly
494 reflecting temperatures that remained relatively low on this river.

495

496 Although limited in being only a rough estimation, parasite abundance was not significantly
497 higher in the large volume samples. This is consistent with previous evidence (the high
498 proportion of false negative samples for 2L samples and qPCR readings close to the LOD)
499 that spore concentrations in the rivers were low at the time of sampling. However, different
500 DNA extraction methods were used for spore samples and river water samples and this may
501 affect the accuracy of our estimates. The handling of the spores collected in the laboratory
502 and those collected as parts of eDNA samples also differed, e.g. the spores from the
503 environment always underwent a filtration procedure which was not the case for laboratory
504 collected spores. This may have further biased the yield of DNA and potentially results in an
505 underestimate of spore numbers deriving from eDNA samples. Sample types should undergo
506 similar treatments in future studies to ensure parallel conversions to spore numbers.
507 Moreover, the fragile nature of malacosporean spores may result in lysis during water
508 filtration and subsequent loss of DNA through the fixed-pore size membranes. Use of glass
509 fibre filters may guard against the latter scenario, as such filters are known to bind free DNA
510 (Nygaard & Hall 1963). Certainly, the abundances of *T. bryosalmonae* inferred by qPCR
511 were much lower than those estimated in similar studies on myxosporean myxozoans. These
512 contrasting abundances may be explained by the more robust nature and greater longevity of
513 spores produced by myxosporeans relative to those of malacosporeans. For example, Hallett
514 & Bartholomew (2006) detected 1 - 20 spores/L of the myxosporean, *C. shasta*, in river water
515 and Griffin et al. (2009) detected 37 - 249 spores/L of the myxosporean, *H. ictaluri*, in pond
516 water. On the other hand, Alama-Bermejo et al. (2013) only detected up to 1 spore of the
517 myxosporean *Ceratonova puntazzi* in 8L of sea water. It is possible that spore abundances of
518 some myxozoans are naturally low. Further investigation is, however, required as we sampled
519 water over a relatively short period of time, at a similar time each day, and did not
520 simultaneously sample bryozoan populations to ascertain spore production.

521

522 ***Caveats and recommendations for future studies***

523 The primers developed for the assay were verified to be specific to the genus
524 *Tetracapsuloides*. Alignments with other putative *Tetracapsuloides* species (Bartošová-
525 Sojková et al. 2014) were inspected, but no DNA isolates were available for testing with the

526 primers developed in this study. Since the primers contain 3 - 6 mismatches to the most
527 closely related *Tetracapsuloides* species (Bartošová-Sojtková et al. 2014), they may also
528 amplify SSU rDNA from these species, especially in the absence of the specific target.
529 Therefore, in environments where the fish fauna is diverse, we recommend post-qPCR
530 sequencing to further verify results. We also recommend examining the melting curve to
531 distinguish between true and false positive samples. Melting curves should produce a sharp
532 peak at ~84.5°C for true positive samples while false positive samples will only amplify
533 primer dimer with a wide peak at ~77.3°C. However, some variation in the template peak
534 temperature, due to either minor pipetting error or nucleotide differences when analysing
535 samples potentially containing different *T. bryosalmonae* strains, should be taken into
536 consideration.

537

538 A potential limitation in the use of environmentally derived DNA to study parasites with
539 multiple host life-cycles is the inability to distinguish between parasite stages released from
540 different hosts. For example, *T. bryosalmonae* DNA detected in our water samples could
541 have four potential sources: (1) Spores that developed and matured in bryozoans; (2) Spores
542 originating from salmonids; (3) Fragments of infected bryozoan colonies present in the water
543 samples; (4) Loose, non-cell associated DNA. Filter papers were closely examined after
544 filtration and no colonies were ever observed. Trapping of loose DNA on fixed cellulose
545 acetate filters is possible as filter papers progressively clog during processing, but this type of
546 filter should predominantly retain only larger particles. Therefore, most of the signal for *T.*
547 *bryosalmonae* is likely to be attributed to spores that were released from bryozoan hosts,
548 particularly since sampling was conducted directly downstream from known infected
549 bryozoan populations and when spore production was expected to be relatively high (Fontes
550 pers. obs). Further work is required to characterise temporal variation in abundances of
551 sporogonic stages in renal tubules of wild fish and the spore loads that are shed from fish.

552

553 There are many potential processes that may impede eDNA-based detection and absolute
554 quantification. Examples include: (1) Loss of DNA if spores rupture during the filtration
555 process; (2) Sample DNA degradation as a result of repetitive freezing/thawing processes;
556 (3) Short viability of *T. bryosalmonae* spores (i.e. 12 – 24 h; Feist et al. 2001, De Kinkelin et
557 al. 2002); and (4) Low DNA extraction efficiency. These biases are largely unavoidable but
558 their impact can be minimised by using appropriate standards, positive and negative controls,

559 well-established and routine sampling methods and post-sampling processes that will enable
560 relative and robust comparisons and maximise information gain. For instance, as mentioned
561 above, lysis of DNA through fixed-pore size membranes could be avoided by using glass
562 fibre filters that bind free DNA. In general, the main sources of uncertainty for water sample-
563 based detection methods for parasites and pathogens stem from spatial and temporal variation
564 in the distribution of target organisms, as we have found here. Repeated sampling, increased
565 replication and larger sample volumes may be required to address these issues.

566

567 The assay developed here provides a tool to resolve parasite abundances over fine time scales
568 and for longer periods. Such studies, for example, may detect seasonal peaks and troughs in
569 the abundance of *T. bryosalmonae*, providing insights as to when transmission is generally
570 achieved in the complex life cycle. Meanwhile, daily variation in estimated spore
571 concentrations in water may provide evidence for spore release entrained to a circadian
572 rhythm, coinciding with fish host activities and increased transmission success. Other
573 applications of our assay include examining how *T. bryosalmonae* abundances change over
574 environmental gradients, with the presence and absence of fish farms sustaining PKD
575 outbreaks, and with hydrological connectivity in river networks.

576

577 **Conclusions**

578

579 We present the first eDNA-based protocol for the detection and quantification of *T.*
580 *bryosalmonae* spores in freshwater samples. Our SYBR[®] Green qPCR assay combined with
581 an IPC provides an easy and rapid method to detect and quantify *T. bryosalmonae*. We have
582 used the assay to characterise variation in spore presence and provide an estimate of
583 abundance in space and time in three river systems. Our new qPCR assay offers a non-
584 destructive means of determining infection risk that may be used to complement traditional
585 monitoring methods.

586

587 **Acknowledgements**

588

589 We are thankful to Darren Butterworth (Trafalgar Fisheries [River Avon]), Robert Starr
590 (Hungerford Fishery [River Dun]), Mrs Pearse and Winchester College (River Itchen) for
591 access to collecting sites. We would also like to thank all field assistants based at the Natural

592 History Museum (Brian Smith, Laetitia Gunton, Alex Gruhl, Graihagh Hardinge, Jahcub
593 Trew) and Nick Taylor (Centre for Environment, Fisheries & Aquatic Sciences [Cefas]) and
594 Chris Williams (Environment Agency [EA]) for their support for research on assessing the
595 risk of PKD. We thank Sascha Hallett (Oregon State University) for initial advice on
596 techniques. The research was funded by the Natural Environment Research Council
597 (NE/019227/1), the EA and Cefas. IF was also funded by the Fundação para a Ciência e a
598 Tecnologia (SFRH/BD/86118/2012) and Pescanova, SA. HH was funded by the Swiss
599 National Science Foundation Sinergia project CRSII3_147649.

600

601 **References**

602

603 Alama-Bermejo G, Šíma R, Raga JA, Holzer AS (2013) Understanding myxozoan infection
604 dynamics in the sea: seasonality and transmission of *Ceratomyxa puntazzi*. Int J
605 Parasitol 43:771-780

606 Audemard C, Ragone Calvo LM, Paynter KT, Reece KS, Bureson EM (2006) Real-time
607 PCR investigation of parasite ecology: in situ determination of oyster parasite
608 *Perkinsus marinus* transmission dynamics in lower Chesapeake Bay. Parasitology
609 132:827-842

610 Bartošová-Sojtková P, Hrabcová M, Pecková H, Patra S, Kodádková A, Jurajda P, Tysl T,
611 Holzer AS (2014) Hidden diversity and evolutionary trends in malacosporean parasites
612 (Cnidaria: Myxozoa) identified using molecular phylogenetics. Int J Parasitol 44:565-
613 577

614 Bates D, Maechler M, Bolker B, Walker S (2013) lme4: Linear mixed-effects models using
615 Eigen and S4. Version: 1.0-4 [software package]

616 Berdal KG, Holst-Jensen A (2001) Roundup Ready® soybean event-specific real-time
617 quantitative PCR assay and estimation of the practical detection and quantification
618 limits in GMO analyses. Eur Food Res Technol 213:432-438

619 Bettge K, Wahli T, Segner H, Schmidt-Posthaus H (2009) Proliferative kidney disease in
620 rainbow trout: time- and temperature-related renal pathology and parasite distribution.
621 Dis Aquat Org 83:67-76

622 Canning EU, Curry A, Feist SW, Longshaw M, Okamura B (2000) A new class and order of
623 myxozoans to accommodate parasites of bryozoans with ultrastructural observations on
624 *Tetracapsula bryosalmonae* (PKX organism). J Eukaryot Microbiol 47:456-468

625 Canning EU, Curry A, Hill SLL, Okamura B (2007) Ultrastructure of *Buddenbrockia allmani*
626 n. sp.(Myxozoa, Malacosporea), a parasite of *Lophopus crystallinus* (Bryozoa,
627 Phylactolaemata). J Eukaryot Microbiol 54:247-262

628 De Kinkelin P, Gay M, Forman S (2002) The persistence of infectivity of *Tetracapsula*
629 *bryosalmonae*-infected water for rainbow trout, *Oncorhynchus mykiss* (Walbaum). J
630 Fish Dis 25:477-482

631 Feist SW, Longshaw M, Canning EU, Okamura B (2001) Induction of proliferative kidney
632 disease (PKD) in rainbow trout *Oncorhynchus mykiss* via the bryozoan *Fredericella*
633 *sultana* infected with *Tetracapsula bryosalmonae*. Dis Aquat Org 45:61-68

634 Feist SW, Morris DJ, Alama-Bermejo G, Holzer AS (2015) Cellular processes in myxozoans.
635 In: Okamura B, Gruhl A, Bartholomew JL (eds) Myxozoan evolution, ecology and
636 development. Springer International Publishing, Cham, Switzerland

637 Ferguson HW, Needham EA (1978) Proliferative kidney disease in rainbow trout *Salmo*
638 *gairdneri* Richardson. J Fish Dis 1:91-108

639 Fiala I, Bartošová-Sojková P, Okamura B, Hartikainen H (2015) Adaptive radiation and
640 evolution within the Myxozoa. In: Okamura B, Gruhl A, Bartholomew JL (eds)
641 Myxozoan evolution, ecology and development. Springer International Publishing,
642 Cham, Switzerland

643 Fontes, I. (2015) Life history, distribution and invertebrate host-parasite interactions of the
644 causative agent of proliferative kidney disease (PKD), *Tetracapsuloides bryosalmonae*.
645 PhD, University of Aberdeen, Aberdeen

646 Fontes I, Hallett S, TorAtle M (2015) Comparative epidemiology of myxozoan diseases. In:
647 Okamura B, Gruhl A, Bartholomew JL (eds) Myxozoan evolution, ecology and
648 development. Springer International Publishing, Cham, Switzerland

649 Foott JS, Stone R, Wiseman E, True K, Nichols K (2007) Longevity of *Ceratomyxa shasta*
650 and *Parvicapsula minibicornis* Actinospore Infectivity in the Klamath River. J Aquat
651 Anim Health 19:77-83

652 Francois P, Pittet D, Bento M, Pepey B, Vaudaux P, Lew D, Schrenzel J (2003) Rapid
653 detection of methicillin-resistant *Staphylococcus aureus* directly from sterile or
654 nonsterile clinical samples by a new molecular assay. J Clin Microbiol 41:254-260

655 Gay M, Okamura B, de Kinkelin P (2001) Evidence that infectious stages of *Tetracapsula*
656 *bryosalmonae* for rainbow trout *Oncorhynchus mykiss* are present throughout the year.
657 Dis Aquat Org 46:31-40

658 Grabner DS, El-Matbouli M (2008) Transmission of *Tetracapsuloides bryosalmonae*
659 (Myxozoa: Malacosporea) to *Fredericella sultana* (Bryozoa: Phylactolaemata) by
660 various fish species. *Dis Aquat Org* 79:133-139

661 Grabner DS, El-Matbouli M (2009) Comparison of the susceptibility of brown trout (*Salmo*
662 *trutta*) and four rainbow trout (*Oncorhynchus mykiss*) strains to the myxozoan
663 *Tetracapsuloides bryosalmonae*, the causative agent of proliferative kidney disease
664 (PKD). *Vet Parasitol* 165:200-206

665 Griffin MJ, Pote LM, Camus AC, Mauel MJ, Greenway TE, Wise DJ (2009) Application of a
666 real-time PCR assay for the detection of *Henneguya ictaluri* in commercial channel
667 catfish ponds. *Dis Aquat Org* 86:223-233

668 Guy RA, Payment P, Krull UJ, Horgen PA (2003) Real-time PCR for quantification of
669 *Giardia* and *Cryptosporidium* in environmental water samples and sewage. *Appl*
670 *Environ Microbiol* 69:5178-5185

671 Hallett SL, Bartholomew JL (2006) Application of a real-time PCR assay to detect and
672 quantify the myxozoan parasite *Ceratomyxa shasta* in river water samples. *Dis Aquat*
673 *Org* 71:109-118

674 Hartikainen H, Fontes I, Okamura B (2013). Parasitism and phenotypic change in colonial
675 hosts. *Parasitology* 140:1403-1412

676 Hartman LJ, Coyne SR, Norwood DA (2005) Development of a novel internal positive
677 control for Taqman[®] based assays. *Mol Cell Probes* 19:51–59

678 Hedrick RP, Baxa DV, De Kinkelin P, Okamura B (2004) Malacosporean-like spores in urine
679 of rainbow trout react with antibody and DNA probes to *Tetracapsuloides*
680 *bryosalmonae*. *Parasitol Res* 92:81-88

681 Hung YW, Remais J (2008) Quantitative detection of *Schistosoma japonicum* cercariae in
682 water by real-time PCR. *PLoS Negl Trop Dis* 2:e337

683 Ishimaru K, Takumi M, Kazunobu T, Sho S (2014) Seasonal monitoring of *Kudoa yasunagai*
684 from sea water and aquaculture water using quantitative PCR. *Dis Aquat Org* 108:45-
685 52

686 Kent ML, Hedrick RP (1986) Development of PKX myxosporean in rainbow trout *Salmo*
687 *gairdneri*. *Dis Aquat Org* 1:169-182

688 Kumar G, Abd-Elfattah A, Saleh M, El-Matbouli M (2013) Fate of *Tetracapsuloides*
689 *bryosalmonae* (Myxozoa) after infection of brown trout *Salmo trutta* and rainbow trout
690 *Oncorhynchus mykiss*. *Dis Aquat Org* 107:9-18

691 McGurk C, Morris DJ, Adams A (2005) Microscopic studies of the link between salmonid
692 proliferative kidney disease (PKD) and bryozoans. *Fish Vet J* 62:71

693 Morris DC, Morris DC, Adams A (2002) Development of improved PCR to prevent false
694 positives and false negatives in the detection of *Tetracapsula bryosalmonae*, the
695 causative agent of proliferative kidney disease. *J Fish Dis* 25:483-490

696 Morris DJ, Adams A (2006) Transmission of *Tetracapsuloides bryosalmonae* (Myxozoa:
697 Malacosporea), the causative organism of salmonid proliferative kidney disease, to the
698 freshwater bryozoan *Fredericella sultana*. *Parasitology* 133:701-709

699 Morris DJ, Adams A (2008) Sporogony of *Tetracapsuloides bryosalmonae* in the brown trout
700 *Salmo trutta* and the role of the tertiary cell during the vertebrate phase of myxozoan
701 life cycles. *Parasitology* 135:1075-1092

702 Nygaard AP, Hall BD (1963) A method for the detection of RNA-DNA complexes. *Biochem*
703 *Biophys Res Commun* 12:98-104

704 Okamura B, Feist SW (2011) Emerging diseases in freshwater systems. *Freshwat Biol*
705 56:627-637

706 Okamura B, Hartikainen H, Schmidt-Posthaus H, Wahli T (2011) Life cycle complexity,
707 environmental change and the emerging status of salmonid proliferative kidney disease.
708 *Freshwat Biol* 56:735-753

709 Opel KL, Chung D, McCord BR (2010) A study of PCR inhibition mechanisms using real
710 time PCR. *J Forensic Sci* 55:25-33

711 Pinheiro J, Bates D, DebRoy S, Sarkar D, Team RC (2014) nlme: linear and nonlinear mixed
712 effects models. Version: 3.1-117 [software package]

713 R Core Team (2014) R: A language and environment for statistical computing. In: R: A
714 language and environment for statistical computing. R Foundation for Statistical
715 Computing, Vienna, Austria

716 Strand DA, Jussila J, Johnsen SI, Viljamaa-Dirks S, Edsman L, Wiik-Nielsen J, Viljugrein H,
717 Engdahl F, Vrålstad T (2014) Detection of crayfish plague spores in large freshwater
718 systems. *J Appl Ecol* 51:544-553

719 Tops S (2004) Ecology, life history and diversity of malacosporeans. PhD, University of
720 Reading, Reading

721 Tops S, Lockwood W, Okamura B (2006) Temperature-driven proliferation of
722 *Tetracapsuloides bryosalmonae* in bryozoan hosts portends salmonid declines. *Dis*
723 *Aquat Org* 70:227-236

- 724 Tops S, Okamura B (2003) Infection of bryozoans by *Tetracapsuloides bryosalmonae* at sites
725 endemic for salmonid proliferative kidney disease. *Dis Aquat Org* 57:221-226
- 726 Wahli T, Bernet D, Steiner PA, Schmidt-Posthaus H (2007) Geographic distribution of
727 *Tetracapsuloides bryosalmonae* infected fish in Swiss rivers: an update. *Aquat Sci*
728 69:3-10
- 729 Zuur AF, Ieno EN, Walker NJ, Saveliev AA, Smith GM (2009) Mixed effects models and
730 extensions in ecology with R, Springer, New York

731 **Figure legends:**

732 **Figure 1.** Location of Roots 1-3 in the Rivers Avon (a), Dun (b) and Itchen (c) along with
733 ordnance survey grid references. Root 3 is in the most downstream position and was where
734 24L samples were taken. (source: OS Street View layer [TIFF geospatial data], Scale
735 1:10,000, Coverage: UK, Ordnance Survey (GB), Using: EDINA Digimap Ordnance Survey
736 Service, <<http://digimap.edina.ac.uk/>>, Downloaded: 20 April 2015, Updated: 26 September
737 2014. © Crown Copyright and Database Right [20/04/2015]. Maps composed in ESRI
738 ArcGIS 10.0.

739

740 **Figure 2.** Results from the *Tetracapsuloides bryosalmonae* sensitivity test of the qPCR
741 assay. A 1:10 serial dilution of two positive samples (Positive sample 1 - Avon 24L, collected
742 downstream from root 3 in the fourth week; Positive sample 2 - Itchen 2L collected
743 downstream from root 3 in the fifth week). The mean quantification cycle (C_q) values and the
744 respective standard errors are presented for each dilution. Each sample was run in duplicate.


745

746 **Figure 3.** Water temperature and flow during the five week period in the Rivers Avon (a),
747 Dun (b) and Itchen (c). Box plots for water flow and temperature include the interquartile
748 range box, the median and the mean (black circles).

749


750 **Figure 4.** Boxplot of *Tetracapsuloides bryosalmonae* spore numbers as estimated by qPCR
751 in 2L water samples collected at three roots and one 24L water sample collected at the most
752 downstream root each week for a period of five weeks in the Rivers Avon, Dun and Itchen.
753 Data for the 2L samples are averaged across the three roots.

754 **Figure 1**




755
756


757 **Figure 2**
758



759



762 **Figure 4**



763

764 **Table 1.** Statistical results on the various analyses (Likelihood Ratio Tests) undertaken
 765 testing the significance of the effect of explanatory variables on parasite presence and
 766 abundance in river water samples. C_q values, spore numbers, water temperature and water
 767 flow variables were mean centered for the statistical analyses.

768

Analysis		Statistical results			
		<i>D</i>	d.f.	P	
Comparison of mean temperature between rivers:	Dun vs. Itchen	0.192	1	0.778	
	Avon vs. Itchen	35.643	1	0.001	
	Avon vs. Dun	30.603	1	0.016	
Effect of roots within a river on:	Water temperature	0.064	6	1	
	Water flow	0.431	6	<0.001	
Effect of sample volume on: (with date included as a random effect)	C_q values	0.913	1	0.339	
	Spore number	0.423	1	0.516	
-		Odds Ratio	χ^2	d.f.	P
Effect of sample volume on likelihood of spore detection		0.11	10.169	1	0.001
Effect of water temperature on likelihood of spore detection		1.60	7.682	1	0.006

769

770 **Online Table 1.** Results from the qPCR primer optimisation using different volume
 771 combinations of each primer (F = forward and R = reverse) using a sample of a single
 772 *Tetracapsuloides bryosalmonae* spore. The results presented are the mean quantification
 773 cycle (C_q) values and respective standard deviation.

F primer volume (μL or μM)	R primer volume (μL or μM)	C_q value \pm SD
0.05	0.05	38.27 ± 0.69
0.05	0.50	33.95 ± 0.81
0.05	1.00	33.75 ± 2.68
0.50	0.05	31.78 ± 2.27
0.50	0.50	29.49 ± 2.37
0.50	1.00	28.54 ± 2.79
1.00	0.05	33.71 ± 3.41
1.00	0.50	29.51 ± 2.11
1.00	1.00	28.15 ± 3.01

774

775 **Online Table 2.** Results from the qPCR inhibition test using an internal positive control
 776 (IPC) using 24L and 2L samples. The results presented are the mean quantification cycle (C_q)
 777 values and respective standard deviation of reactions without eDNA (IPC control) and with
 778 eDNA spiked (24L and 2L samples).

779

780

781

782

783

784

785

786

787

788

789

790

791

792

793

794

795

Sample		C_q value \pm SD
IPC control		16.34 \pm 0.407
24L	Avon Root 3 Week 1	16.80 \pm 0.262
	Dun Root 3 Week 4	16.54 \pm 0.012
	Itchen Root 3 Week 3	16.81 \pm 0.172
2L	Avon Root 3 Week 1	18.27 \pm 0.289
	Dun Root 2 Week 2	18.70 \pm 0.525
	Dun Root 2 Week 3	18.23 \pm 0.004
	Itchen Root 2 Week 3	16.83 \pm 0.078

796 **Online Table 3.** Results from the *Tetracapsuloides bryosalmonae* qPCR sensitivity test. A
 797 1:10 serial dilution of two positive samples (the three negative samples tested were negative
 798 at any dilution) was tested. The results presented are the mean quantification cycle (C_q)
 799 values and respective standard deviation in PCR reactions with and without the addition of
 800 bovine serum albumin (BSA). There was no significant difference between C_q values
 801 obtained with and without BSA (Paired t-test: $t = -0.837$, d.f. = 5, $P = 0.441$).

Sample	Dilution factor	C_q value \pm SD	
		No BSA	BSA
Itchen 2L Root 3 Week 5	0	21.81 \pm 7.88	21.12 \pm 0.24
	1:10	25.88 \pm 0.18	25.67 \pm 0.34
	1:100	31.00 \pm 0.20	30.76 \pm 0.92
	1:1,000	-	-
	1:10,000	-	30.58 \pm 0.87
	1:100,000	-	-
Avon 24L Root 3 Week 4	0	26.13 \pm 0.12	26.82 \pm 0.09
	1:10	27.46 \pm 0.77	27.64 \pm 0.16
	1:100	-	30.38 \pm 1.23
	1:1,000	-	31.29 \pm 0.03
	1:10,000	-	-
	1:100,000	-	-

802

803 **Online Table 4.** Inter-assay variation (reproducibility) of the *Tetracapsuloides bryosalmonae*
804 assay using all seven reference standard points. Calculated mean concentration, standard
805 deviation (SD) and coefficient of variation (CV) are given (n = 4 runs).
806

Expected standard concentration (mol/L)	Calculated mean concentration (mol/L)	SD	CV (%)
1e-12	1.14e-12	9.56e-14	8.40
1e-13	1.12e-13	1.07e-14	9.60
1e-14	8.92e-15	5.28e-16	5.91
1e-15	7.87e-16	4.15e-17	5.28
1e-16	1.02e-16	9.03e-18	8.88
1e-17	9.25e-18	9.28e-19	10.03
1e-18	1.26e-18	1.11e-19	8.78

807 **Online Table 5.** *Tetracapsuloides bryosalmonae* abundance according to sampling method,
 808 river, root and trip. Abundance is presented as mean quantification cycle (C_q) values and
 809 respective standard deviation.

Sampling method	River	Root	C_q value \pm SD				
			Week 1	Week 2	Week 3	Week 4	Week 5
2L	Avon	1	-	-	29.21 \pm 0.48	-	-
		2	29.80 \pm 0.18	-	29.35 \pm 0.16	-	30.22 \pm 0.13
		3	-	30.58 \pm 0.61	29.51 \pm 1.22	29.97 \pm 0.51	-
	Dun	1	-	-	-	28.27 \pm 0.30	-
		2	-	30.50 \pm 0.59	30.54 \pm 0.05	-	-
		3	29.63 \pm 0.32	-	-	-	-
	Itchen	1	-	-	-	-	30.27 \pm 0.37
		2	-	-	-	-	-
		3	-	-	-	-	25.63 \pm 0.27
24L	Avon	3	29.38 \pm 0.50	27.51 \pm 0.37	29.03 \pm 0.42	26.17 \pm 0.16	29.35 \pm 0.53
	Dun	3	-	30.62 \pm 0.05	30.13 \pm 0.35	29.20 \pm 0.13	28.28 \pm 0.64
	Itchen	3	-	-	-	30.75 \pm 0.61	28.48 \pm 0.43

810

811 **Supplementary figure legends:**

812

813 **Online Figure 1.** Standard curve derived from a 1:10 serial dilution of an 18S

814 *Tetracapsuloides bryosalmonae* DNA template showing parasite DNA concentration vs.

815 quantification cycle (C_q) value. Each standard was run in duplicate. The curve is significantly

816 linear over seven references from $1e-18$ to $1e-12$ mol/L.

817 Online Figure 1

